


AD A110571



Diurnal Changes in Rectal and Body Surface Temperatures of Conscious, Chair-Restrained Rhesus Macaques

C. T. Liu, PhD; R. P. Sanders; V. W. Robbins, DVM

SUMMARY

A system has been established for continuous recording of rectal and body surface temperatures in conscious, chair-restrained rhesus macaques. Diurnal changes in rectal temperature and body surface temperatures of head, back, abdomen, and inner thigh were recorded for 24 hours under conditions of intermittent or constant light. When lights were on for 8 hours (0800 to 1600 hours) and off for 16 hours, maximal rectal temperatures (37.6 ± 0.3 C) were observed at 0900 hours, whereas minimal rectal temperatures (36.7 ± 0.4 C) were demonstrated between 0200 and 0400 hours. Mean rectal temperatures varied 0.9 C during a 24-hour period. A trend of diurnal changes in body surface temperatures was also observed with low surface temperatures recorded between 0400 and 0800 hours. The surface temperatures (33.8 to 36.5 C) were markedly lower than rectal temperatures (36.7 to 37.6 C) under

the same experimental conditions. When macaques were subjected to constant light, high rectal temperatures were recorded between 0900 and 1700 hours and low rectal temperatures were observed between 2400 and 0300 hours with a mean variation of 0.7 C (36.3 ± 0.15 to 37.0 ± 0.2). Diurnal body surface temperature changes during constant light were less pronounced than those of rectal temperatures. Furthermore, low surface temperatures of abdomen and inner thigh recorded between 0200 and 0800 hours were similar to those obtained with macaques exposed to the light-dark condition.

In a review of the literature, few studies reported normal values of rectal and body surface temperatures in rhesus macaques. Comprehensive information on diurnal rectal and body surface temperature changes in the nonhuman primate was also lacking. Erikson¹ reported diurnal vaginal temperatures for 12 hours in 9 unrestrained rhesus macaques kept in a room with constant light for 8 hours and darkness for 16 hours. Vaginal temperatures were measured twice with a clinical thermometer during the period between 1000 and 2200 hours; the interval between tempera-

ture measurements was 3.5 to 5.0 hours. The highest temperature was recorded in early afternoon and the lowest at 2200 hours. The diurnal temperature variation ranged from 0.72 to 1.66 C. Diurnal changes in colonic temperature of restrained squirrel monkeys were determined with a thermistor.² Squirrel monkeys displayed a prominent circadian rhythm in colonic temperature, when exposed each 24 hours to an equal distribution of light-dark cycles. Colonic temperature was also maintained during 6 hours of mild cold exposure (20 C) at all circadian phases. However, in constant light, the squirrel monkey's ability to regulate colonic temperature against similar cold exposure was impaired.

In a classic work, Galbraith and Simpson³ measured the axillary temperature with clinical thermometers for recording diurnal changes in monkeys (species unspecified). When monkeys were kept continuously in the light, there was no regular diurnal wave. The regular diurnal wave was observed only when monkeys were kept in 12 hours of darkness and 12 hours of constant light. The effect of light on diurnal changes in deep body temperature was also demonstrated in caged capuchin monkeys (*Cebus* sp.).⁴ In a study with rhesus macaques, a rhythm of subcutaneous temperature changes was measured by thermistor surgically implanted under the dorsal skin of the neck.⁵ The rhythm was

Received for publication Sept 2, 1980.
From the US Army Medical Research Institute of Infectious Diseases, Fort Detrick, Frederick, MD 21701.

A portion of this work was presented at the Federation of American Societies for Experimental Biology, April 13-18, 1980, Anaheim, Calif. and published as an abstract (*Fed Proc* 39: 989, 1980).

The authors thank Captain D. Foren for writing the computer program for recording of rectal temperatures, Mr. S. Tobery for technical assistance, Mrs. Phebe W. Summers for editorial assistance, and Mrs. Barbara W. Kline for secretarial aid.

influenced by both the lighting and the feeding schedules. Furthermore, Rohles et al⁶ measured the mean body surface temperature at the groin and calf of a chimpanzee, but no detailed procedures were given. Maximal surface temperatures were recorded between 1600 and 2400 hours, indicating a possible diurnal surface temperature variation.

Since the rhesus macaque is commonly used as a primate model for the study of human fever,⁷ toxemia,^{8,9} and infectious diseases,¹⁰⁻¹² simple techniques for measuring rectal and body surface temperatures are essential for performing these experiments. The purposes of the present study were (1) to develop a noninvasive system for obtaining long-term (> 48 hours) continuous recordings of rectal and body surface temperatures of restrained macaques and (2) to establish baseline diurnal rectal and body surface temperature values of conscious, chair-restrained macaques.

Materials and Methods

Experimental Design—Eleven apparently healthy rhesus macaques weighing 4.2 to 10.1 kg were allocated into 2 groups. Group I (n = 5) was subjected to 8 hours' light (0800 to 1600 hours), followed by 16 hours' darkness in a 24-hour period. Group II (n = 6) was exposed to constant light. All macaques were chair-restrained in an isolation room,^a with constant temperature ($23 \pm 1^\circ\text{C}$) and relative humidity ($20 \pm 1\%$) and fed with monkey chow^b at 0900 and 1600 hours. Water was provided ad libitum. Temperature recordings were started approximately 24 hours after chair-restrained procedure.

Preparation of Thermocouple Electrodes—Fabrication of thermocouple electrodes for continuous measurements of body surface temperatures consisted of stripping 1 cm of insulation from the ends of a 3.0 to 3.6 m length of thermocouple wire.^c The exposed strands were twisted together and soldered to a 2-cm diameter brass disk 0.13 mm thick. The soldered junction was strengthened by embedding the disk and an adjacent 5 to 10 mm of wire in plastic cement.^d This was done by forming a mold around the disk and wire with modeling clay. Plastic powder and liquid were mixed and poured into the mold to form a 2 to 3 mm layer over the brass disk (Fig 1). After hardening, the disk was removed from the mold and the




Fig 1—Preparation of the body surface electrode.

rough edges of the plastic were filed smooth. Four or more holes were drilled around the periphery of the disk to facilitate suturing. The other end of the wire was attached to the terminals of a plug assembly^e which linked the electrode to a potentiometer.^f The potentiometer was calibrated with the electrode immersed in a constant temperature waterbath.^g

Measurements of Rectal and Body Surface Temperature—Approximately 24 hours before temperature recordings, the macaques were sedated with ketamine (10 mg/kg). Four thermocouple electrodes were sutured to the occipital region of the head, lower abdomen, back (thoracic region), and inner thigh, respectively. Skin temperatures from the 4 different sites of the conscious, chair-restrained macaques were recorded continuously on a potentiometer.^h The rectal temperature was measured with a stainless steel rectal probe,ⁱ lubricated with sterile jelly, and inserted 3 to 4 cm into the rectum. The rectal probe, secured to the tail with adhesive tape, was connected to a telether-

meter.^j The output signals were fed into a minicomputer^k via a universal (DC) preamplifier of a recorder^l (Fig 2). A computer program (Fig 3) for printing out rectal temperatures from signals of the recorder was incorporated into the recording system. The calibration procedures were identical, as described for thermocouple electrodes. All temperatures were recorded continuously for 48 hours. Data from a complete 24-hour cycle (2400 to 2400 hours) were analyzed to obtain mean values and SEM.

Results

Rectal Temperature—Diurnal changes in rectal temperature were observed in the conscious, chair-restrained rhesus macaques exposed to constant light for 24 hours or with 8 hours' light exposure (0800 to 1600 hours) (Fig 4). With lights on for 8 hours and off for 16 hours (complete darkness), maximal rectal temperatures ($37.6 \pm 0.3^\circ\text{C}$) were recorded at 0900 hours and decreased thereafter. Minimal rectal temperatures ($36.7 \pm 0.4^\circ\text{C}$) were demonstrated between 0200 to 0400 hours. Mean rectal tem-

^a Halston Purina Co, St Louis, Mo.
^b 28 AWG, 2-conductor, Revere Corp, Wallingford, Conn.
^c Cranioplastic Cement, Plastic Products Co, Roanoke, Va.

^d Model 7313, Honeywell Industrial Division, Fort Washington, Pa.
^e Elektronik 111/112, Honeywell Industrial Division, Fort Washington, Pa.
^f Polyscience waterbath (Model J4-9621), American Instrument Co, Silver Springs, Md.
^g Rectal probes (Model 403), Yellow Springs Instrument Co, Yellow Springs, Ohio.
^h Telethermometer (Model 44TA), Yellow Springs Instrument Co, Yellow Springs, Ohio.

ⁱ Minicomputer—Digital pdp 11/34, Digital Equipment Corp, Maynard, Mass.
^j Recorder—Model Mark 200, Brush Instruments Division, Cleveland, Ohio.




Fig 2—Schema for recording rectal and body surface temperatures in conscious chair-restrained rhesus macaques.

peratures varied 0.9 C during a 24-hour period. When macaques were subjected to constant light, high rectal temperatures were recorded between 0900 and 1700 hours and decreased thereafter. Low rectal temperatures were observed between 2400 and 0300 hours. The diurnal rectal temperatures varied from 36.3 ± 0.15 C to 37.0 ± 0.2 C; a change of 0.7 C.

Body Surface Temperature—Under conditions of 8 hours' light and 16 hours' darkness, a trend of diurnal changes in surface temperatures was observed on the head, back, abdomen, and inner thigh (Fig 5). In general, low surface temperatures were recorded between 0400 and 0800 hours, and high surface temperatures, between 1000 to 1500 hours. The surface temperatures (33.8 to 36.4 C) were markedly lower than rectal temperatures (36.7 to 37.6 C) under the same experimental conditions. With lights on at all times, diurnal surface temperature changes became difficult to identify except at the sites of the abdomen and the inner thigh. Low surface temperatures were recorded between 0200 and 0800 hours.

Table 1 summarizes the results of maximal and minimal temperature changes of rectal and individual surface temperature changes of rhesus macaques within 24 hours. The time or duration for recording maximal and minimal temperatures are also indicated. Constant light impaired diurnal changes of rectal and surface temperatures in general, especially for the surface of the head and back.

Discussion

Diurnal temperature change is a normal phenomenon of biological variation during a nearly 24-hour period. Physiologic and biochemical aspects of circadian rhythm and neural mechanisms for the regulation of these periodic changes have been reviewed by Menaker,¹³ Aschoff and Weber,¹⁴ and Rusak and Zucker.¹⁵ Effects of food and light on modification of circadian rhythm were reported by Iampietro et al¹⁶ and Fuller et al.²

Elizondo¹⁷ indicated that the "normal" core body temperature of many mammalian species ranged from 36 to 39 C. The reported values of "body" temperature of normal rhesus ma-

caques^{18,19} fall within these limits. "Body" temperatures in primates have been determined from different sites including back muscle (longissimus dorsi),^{18,19} vagina,¹ axilla,³ and esophagus²⁰ (under anesthesia). Baseline rectal temperatures obtained from the present study ranged from 36 to 37 C under constant light. These rectal temperature values are similar to human body temperature (37 C), but lower than those of reported values obtained from back muscles for rhesus macaques.^{18,19} The difference in normal body temperature of the same primate species may be due to different techniques used. For example, when body temperature is measured from the back muscle with thermocouple wires as demonstrated by others,^{18,19} the traumatized or injured muscle may produce local vasodilation²¹ and increase the temperature. Furthermore, emotional stress resulting from human handling of the conscious macaque by inserting a probe or clinical thermometer into the rectum may also increase the rectal temperature. However, in the present study, we attempted to eliminate these disturbing factors, so that more

```

FILE 11: TEMP2.FIN
DIMENSION TEMP(4),IT(3),TIM(2),ATEMP(29)
INTEGER IBUF(4,2),IBUF(40),IBUFNO,ADNUM(30)
IBFNT=0      !IBFNT WILL TELL NUMBER OF PASSES THROUGH
TYPE *, ' IS SAMPLE TIME TO BE MEASURED IN SECONDS OR MINUTES S,M'
ACCEPT 45,ALPHA
45  FORMAT(A1)
    IF(ALPHA.EQ.'M')GO TO 55
    TYPE *, ' ENTER CHANNEL # AND SECONDS BETWEEN SAMPLES'
    ACCEPT 40,ICHAN,ITIME
    FORMAT(2I5)
    IPRSET=-100*ITIME
    CALL CLOCKA(5,IPRSET)      !CLOCK RATE SET AT 100 HZ
    GO TO 35
55  TYPE *, ' ENTER CHANNEL # AND MINUTES BETWEEN SAMPLES'
    ACCEPT 41,ICHAN,ITIME
    FORMAT(2I5)
    CALL CLOCKB(5,-100,2,IND)
    IF(IND.EQ.1) GO TO 44
    TYPE *, ' ERROR IN LOADING CLOCKB'
44  CONTINUE
    IPRSET=-60*ITIME
    CALL CLOCKA(0,IPRSET)
35  TYPE *, ' ENTER NUMBER OF SAMPLES TO BE TAKEN'
    TYPE *, ' # OF SAMPLES MUST BE A MULTIPLE OF 4 OR IT WILL BE TRUNCATED'
    ACCEPT 50,ISAMPL  !FOR PRESENT MULTIPLE OF 4
50  FORMAT(I5)
    CALL CLREF(30)
    CALL SETIBF(IBUF,,,BUF(1,1),BUF(1,2))
    CALL RLSBUF(IBUF,,,0,1)
    CALL ADSWF(IBUF,4,0,64,1,30,0,ICHAN,1)
75  CALL DATE(DT)
    CALL TIME(TIM)
    TYPE 12
12  FORMAT(1H,/, ' DATE',9X,' TIME')
    TYPE 10,DT,TIM
10  FORMAT(1H,3A4,4X,2A4)
    IBUFNO=IWTRUF(IBUF,30) + 1
    IF(IBUFNO.LT.1) GO TO 99  !STOP PROGRAM
    TYPE 15
15  FORMAT(1H,29X,' TEMPERATURE')
C ***** DECISION ROUTINE TO DETERMINE TEMPERATURE VALUES *****
DO 300 I=1,4
    IF(BUF(I,IBUFNO).LE.ADNUM(1)) GO TO 250
DO 200 J=1,29
    K=J+1
    IF(BUF(I,IBUFNO).GT.ADNUM(1).AND.(BUF(I,IBUFNO).LE.ADNUM(K))) GO TO 250
200  CONTINUE
C  OVERFLOW
    TEMP(I)=99.
    GO TO 500
220  TEMP(I)=ATEMP(I)
    GO TO 500
C  UNDERFLOW
250  TEMP(I)=0.
C *****
500  WRITE(5,100)BUF(I,IBUFNO),TEMP(I)
100  FORMAT(1H,13X,18,5X,F10,2)
300  CONTINUE
    NBUF=IBUFNO-1
    CALL RLSBUF(IBUF,,,NBUF)
    IBFNT=IBFNT+1
    IF(IBFNT.EQ.ISAMPL/4) GO TO 99  !STOP
    GO TO 25
99  CALL STEPSW(TRUE,0)
    DATA ADNUM/2161,2192,2200,2240,2255,2276,2301,2322,2338,
    1,2362,2393,2416,2433,2453,2478,2499,2416,2536,2558,2580,2601,
    2,2623,2645,2665,2683,2700,2733,2758,2781,2804/
    DATA ATEMP/35.,45,25,35,5,35,75,36.,36,25,36,5,36,25,37.,
    137,25,37,5,37,25,38.,38,25,38,5,38,25,39.,39,25,39,5,
    239,25,40.,40,25,40,5,40,25,41.,41,25,41,5,41,25,42,7
    STOP
    END

```

Fig 3—Computer program for recording rectal temperatures.




Fig 4—Diurnal changes in rectal temperatures of conscious, chair-restrained rhesus macaques.

TABLE 1—Summary of Results

| Location | Diurnal rectal and body surface temperature changes | | | Time of maximal and minimal temperature changes (hours of day) | |
|---|---|---------------------|--|--|------------|
| | Maximal temperature | Minimal temperature | Maximal temperature change during 24 hours | Maximum | Minimum |
| 8 HOURS' LIGHT + 16 HOURS' DARKNESS (n = 5) | | | | | |
| Rectal | 37.6 ± 0.3 | 36.7 ± 0.4 | 0.9 | 0900 | 0200-0400 |
| Body surface | | | | | |
| Head | 36.2 ± 0.6 | 35.1 ± 0.3 | 1.1 | | |
| Abdomen | 34.9 ± 1.0 | 33.8 ± 0.8 | 1.1 | | |
| Back | 35.8 ± 0.7 | 34.3 ± 0.6 | 1.5 | 1000-1500 | 0400-0800 |
| Inner thigh | 36.4 ± 0.3 | 35.1 ± 0.5 | 1.3 | | |
| 24 HOURS' CONSTANT LIGHT (n = 6) | | | | | |
| Rectal | 37.0 ± 0.2 | 36.3 ± 0.2 | 0.7 | 0900-1700 | 2400-0300 |
| Body surface | | | | | |
| Head | 35.3 ± 0.8 | 34.5 ± 0.8 | 0.8 | | |
| Abdomen | 36.8 ± 0.7 | 34.8 ± 0.8 | 1.0 | Not clearly evident | |
| Back | 35.9 ± 0.7 | 34.8 ± 0.8 | 1.1 | | |
| Inner thigh | 36.3 ± 0.8 | 35.0 ± 0.9 | 1.3 | | 0200-0800* |

* Only for abdominal and inner thigh surface temperatures.

closely realistic values of rectal temperatures were obtained. Diurnal changes of rectal and body surface temperatures in conscious chair-restrained rhesus macaques varied approximately 1 C during a 24-hour pe-

riod. Surface temperatures were lower than the rectal temperature but the timing for reaching maximal and minimal values was different between surface and rectal temperatures. Since surface temperature changes might be

associated with cutaneous vasoconstriction and vasodilation for body temperature regulation under various environmental light conditions, this may explain why diurnal skin changes were less pronounced than changes of rectal temperatures.

Baker et al²² demonstrated variability of body surface temperatures at different sites of conscious pigtail macaques (*Macaca nemestrina*). Different base-line values for tail, foot, and mean surface temperatures were also observed in the squirrel monkey (*Saimiri sciureus*) by Adair and Adams.²³ Although surface temperatures of the inner thigh^{24,25} and ear of rhesus macaques have been reported, base-line thigh surface temperatures (37.2 to 37.9 C) are markedly higher than those values obtained from the same site in the present study (36.4 ± 0.3 C at 0900 hours). This discrepancy might be due to different techniques in measuring surface temperature and different experimental conditions. Higher thigh surface temperatures recorded by other investigators might be the result of surgical operation, in which a durable thermode implant was placed in the hypothalamus.²⁵ Furthermore, the complicated experimental design and restraint of the macaque may also provide additional stress to alter surface temperature.

The selection of a rigid stainless steel rectal probe was based on the evidence that chair-restrained macaques were not able to push the probe out of the anus as easily as a flexible, soft probe. However, the main disadvantage for using the rigid rectal probe was the possibility of damaging the rectum when the macaque was too active or could not be easily adapted to chairing. Under this circumstance, the use of a soft rectal probe is recommended, providing that the segment of the probe wire outside of the anus is supported by a rigid rod and fixed firmly to the macaque's tail with adhesive tape. The temperature recording system described here not only provides less stress on the animal, but also operates simply and automatically for data processing. Because rhesus macaques reveal certain diurnal changes of rectal and body surface temperatures, the timing for daily measurements of temperatures should be consistent to avoid errors. Under normal circumstances, the diurnal temperature variation of a rhesus macaque may be approximately 1 C.




Fig 5—Diurnal changes in body surface (head, inner thigh, abdomen, and back) temperatures of conscious chair-restrained rhesus macaques.

However, when fever is induced, diurnal temperature changes are difficult to predict. Thus, a daily continuous recording of rectal and body surface temperatures appears to be the best approach for correct and accurate interpretations of data.

References

1. Erikson LB: Diurnal temperature variation in the rhesus monkey under normal and experimental conditions. *Nature (London)* 186: 83-84, 1960.
2. Fuller CA, Sulzman RM, Moore-Ede MC: Effective thermoregulation in primates depends upon internal circadian synchronization. *Comp Biochem Physiol* 63A:207-212, 1979.
3. Galbraith JJ, Simpson S: Conditions influencing the diurnal wave in the temperature of the monkey. *Proc Physiol Soc (London)* 20-22, 1903.
4. Winget CM, Card DH, Hetherington NB: Circadian oscillations of deep-body temperature and heart rate in a primate (*Cebus albifrons*). *Aerosp Med* 39:350-353, 1968.
5. Ternes JW, Farrer DN: A circadian temperature rhythm in the rhesus monkey. Report No. ARL-TR-66-21. 6571st Aeromedical Research Laboratory, Holloman Air Force Base, New Mexico, 1966.
6. Rohles FH, Reynolds H, Brown C: Diurnal temperature changes in a chimpanzee during a 14-day restraint test. Report No. ARL-TDR-63-2. 6571st Aeromedical Research Laboratory, Holloman Air Force Base, New Mexico, 1963.
7. Perlow M, Dinarello CA, Wolff SM: A primate model for the study of human fever. *J Infect Dis* 132:157-164, 1975.
8. Liu CT, DeLauter RD, Faulkner RT: Cardiovascular and hepatic responses of rhesus macaques to staphylococcal enterotoxin B. *Am J Vet Res* 38:1849-1854, 1977.
9. Liu CT, DeLauter RD, Griffin MJ, et al: Effects of staphylococcal enterotoxin B on functional and biochemical changes of the lung in rhesus monkeys. *Toxicon* 16:543-550, 1978.
10. Kastello MD, Spertzel RO: The rhesus monkey as a model for the study of infectious disease. *Am J Phys Anthropol* 38:501-504, 1973.
11. Liu CT, Hilmas DE, Griffin MJ, et al: Alterations of body fluid compartments and distribution of tissue water and electrolytes in

rhinus monkeys with Rocky Mountain spotted fever. *J Infect Dis* 138:42-48, 1978.

12. Liu CT, Sanders RP, Hadick CL, et al: Effects of intravenous injection of leukocytic endogenous mediator on cardiohepatic functions in rhesus macaques. *Am J Vet Res* 40: 1035-1039, 1979.

13. Menaker M: Physiological and biochemical aspects of circadian rhythms. Introductory remarks. *Fed Proc* 35:2325, 1976.

14. Aschoff J, Weber R: Human circadian rhythms: A multioscillatory system. *Fed Proc* 35: 2326-2332, 1976.

15. Rusak B, Zucker I: Neural regulation of circadian rhythms. *Physiol Rev* 59:449-526, 1979.

16. Iampietro PF, Buskirk ER, Bass DE,

et al: Effect of food, climate and exercise on rectal temperature during the day. *J Appl Physiol* 11:349-352, 1957.

17. Elizondo R: Temperature regulation in primates. *Int Rev Physiol Environ Physiol* 15: 71-118, 1977.

18. Bilbrey GL, Beisel WR: Depression of free water clearance during pneumococcal bacteremia. *Ann Surg* 177:112-120, 1973.

19. Boucher JH, Hilmas DE, Liu CT, et al: Myocardial depression during *Diplococcus pneumoniae* infection in monkeys. *Proc Soc Exp Biol Med* 145:112-116, 1974.

20. Kopf GS, Mirvis DM, Myers RE: Central nervous system tolerance to cardiac arrest during profound hypothermia. *J Surg Res* 18: 29-34, 1975.

21. Liu CT: Circulatory response to muscle trauma following denervation and hemorrhage in dogs. *J Trauma* 8:75-83, 1968.

22. Baker MA, Cronin MJ, Mountjoy DG: Variability of skin temperature in the waking monkey. *Am J Physiol* 230:449-455, 1976.

23. Adair ER, Adams BW: Microwaves induce peripheral vasodilatation in squirrel monkey. *Science* 207:1381-1383, 1980.

24. Johnson GS, Elizondo RS: Thermoregulation in *Macaca mulatta*: A thermal balance study. *J Appl Physiol* 46:268-277, 1979.

25. Smiles KA, Elizondo RS, Barney CC: Sweating responses during changes of hypothalamic temperature in the rhesus monkey. *J Appl Physiol* 40:653-657, 1976.

| | |
|--------------------|-------|
| Accession For | |
| NTIS GRA&I | |
| DTIC TAB | |
| Unannounced | |
| Justification | |
| By | |
| Distribution | |
| Availability Codes | |
| Avail and/or | |
| Dist | Specs |
| A 21 | |

